

Maximum Strength Mucinex[®] DM

Drug Facts

| | |
|--|----------------|
| Active ingredients (in each extended-release bi-layer tablet) | Purpose |
|--|----------------|

| | |
|---------------------------------|-------------------|
| Dextromethorphan HBr 60 mg..... | Cough suppressant |
| Guaifenesin 1200 mg..... | Expectorant |

- Uses**
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
 - temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Warnings

- Do not use**
- for children under 12 years of age
 - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

When using this product

- do not use more than directed

Drug Facts (continued)

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older:
1 tablet every 12 hours; not more than 2 tablets in 24 hours
- children under 12 years of age: do not use

Other information

- tamper evident: do not use if carton is open or if seal on bottle printed "SEALED for YOUR PROTECTION" is broken or missing
- store at 20-25°C (68-77°F)

Inactive ingredients carbomer 934P, NF; D&C yellow #10 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF