

# Mucinex® D

## Drug Facts

<b>Active ingredients</b> <b>(in each extended-release</b> <b>bi-layer tablet)</b>	<b>Purpose</b>
--	----------------

Guaifenesin 600 mg.....	Expectorant
Pseudoephedrine HCl 60 mg.....	Nasal Decongestant

### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves nasal congestion due to:
  - common cold
  - hay fever
  - upper respiratory allergies
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- temporarily relieves sinus congestion and pressure

### Warnings

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

#### When using this product

- do not use more than directed

## Drug Facts (continued)

### Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days, come back or occur with a fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for timing of meals
- adults and children 12 years and older: 2 tablets every 12 hours; not more than 4 tablets in 24 hours
- children under 12 years of age: do not use

### Other information

- tamper evident: do not use if carton is open or if printed seal on blister is broken or missing
- store at 20-25°C (68-77°F)

**Inactive ingredients** carbomer 934P, NF; FD&C yellow #6 aluminum lake; hypromellose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF